

# Electroencephalographic Identifiers of Reading Abilities in Turkish Language

Günet Eroğlu

Faculty of Engineering and Natural Sciences  
Sabanci University, İstanbul, Turkey  
gunet.eroglu@healthmobilesoftware.com

Müjdat Çetin

Faculty of Engineering and Natural Sciences  
Sabanci University, İstanbul, Turkey  
Dept. of Electrical and Computer Eng., University of Rochester, Rochester, NY, USA  
mccetin@sabanciuniv.edu

Selim Balcisoy

Faculty of Engineering and Natural Sciences  
Sabanci University, İstanbul, Turkey  
baltisoy@sabanciuniv.edu

**Abstract**—Fluent reading requires learning the print knowledge of alphabet symbols (letters), rapid automatic naming and phonological awareness skills. In this study, electroencephalographic brain signals of 17 subjects were measured with an eMotiv EPOC+ headset before, during and after a computer-based training session. For the training, distorted letter prints were created by rotating Turkish letters 180 degrees along the y-axis. Using these distorted letters two different texts, each 150 words in length, were created. Subjects were asked to read these texts before and after the training session. We investigated whether there is an improvement in reading speed and a decrease in number of errors due to the computer-based training and whether we can correlate the success of training with any characteristic of any EEG brain signals. Based on our analysis of the EEG data collected throughout the experiment, we observed that the frequency modulation across resting states in the Theta at the Broca Area (F7 and FC5) predicts individual reading performance measures. Even though there exist a variety of studies indicating a relation of Theta band power and learning performance, EEG measurements with eMotiv EPOC+ had not been previously reported with a Turkish alphabet learning task.

**Keywords**—EEG signal processing, multi sensory learning.

## I. INTRODUCTION

Reading, writing and arithmetic abilities are important for academic success. University students pass many exams throughout their school lives by the time they reach university education. However, in spite of the fact that their IQ is normal or above normal, some of these students, suffer from specific learning disabilities. If the Wechsler Intelligence Scale for Children (WISC-R) test shows there is a big discrepancy between verbal IQ and performance IQ, this may point to specific learning disability (SLD). However, there are many specific learning disability cases which cannot be shown by such tests

alone [11]. The findings in the article suggest that IQ tests are not particularly useful for determining learning disabilities. Instead achievement (reading, arithmetic, writing) test scores are more useful diagnostic tools. Learning to read starts with learning the names of the alphabet symbols (rapid automatic naming) and their related sounds (phonological awareness) [1]. The relation between an alphabet symbol and its sound should be unique in order to retrieve the information correctly later on when needed during reading. Reading speed is based on how a person easily retrieves the information about the symbol and its sound(phonics). This reading process involves both short-term and long-term memory. Both rapid automatic naming and phonological awareness skills have been strongly predictive of individual differences in reading. The performance measure for good reading is, therefore, the reading speed (number of words read in a minute) and the number of errors made during reading. Reading comprehension (which is to understand what is read) develops in the brain after the reading speed reaches a certain level and the vocabulary is extended enough, which requires a variety of oral-language skills [2], [3]. Although IQ is normal or above normal, 5% of people in many societies may not perform well in reading, writing and/or arithmetic, and/or in learning in general [4], [7]. There is a great deal of research about specific learning disabilities in the literature which show that specific learning disabilities are mostly developmental [12], [13], [24]. The research conducted in this area suggests an abnormal lateralization of prefrontal attentional control processes [22]. This finding is in accordance with the (scarce) evidence that reading disabilities involve a deviant structural asymmetry of the frontal lobe [23]. The recent fMRI studies in neurotypical adults implicate corticostriatal and hippocampal systems in language learning [27]. In the literature, it has been further shown that higher levels of slow brain waves in the Broca area (F7 and FC5) in the left hemisphere have a high correlation with linguistic learning disabilities [5], [6], [10], [21], [25], [26]. These studies have been completed

using laboratory EEG equipment with 32 or 64 electrodes. In our study, we measured the EEG signals of subjects with a lightweight eMotiv EPOC+ headset with 14 electrodes before, during and after a learning task. The Emotiv EPOC+ headset (<http://emotiv.com/>) has become one of the leading pieces of BCI equipments available today as a result of its low cost and features, including 14 EEG channels plus two references, inertial sensors, wireless connectivity, the ability to provide raw EEG data as well as averaged EEG data, ease of use and attractiveness for children and adolescents. Although its use for research purposes is being questioned by many researchers, several researches have demonstrated that eMotiv EPOC+ headset captures the EEG data quite accurately [?], [8], [19], [20]. In order to collect data from many children, eMotiv EPOC+ provides a user friendly and ergonomic interface and can integrate with both desktop and mobile phone applications and be used at home. In the near future, with this new technology, it can be possible to apply neurofeedback to young children at home without disturbing them much and without lowering their self-esteem. Turkish is an ortographic language, meaning that the words are read as they are written. This makes the reading easier for many people, yet there are learning disabilities in Turkey. There has been little research [14]–[18] conducted about specific learning disabilities in Turkey and to our knowledge, there is no research conducted about Turkish language which correlates learning disabilities with EEG data. The contribution of this paper is an experimental study that explores neural correlates of reading abilities in Turkish based on EEG data collected by an eMotiv EPOC+ headset. Our experimental analysis of the collected data show that higher slow brain waves in the Broca area (F7 and FC5) in the left hemisphere have a high correlation with linguistic learning disabilities.

## II. MATERIALS & METHODS

### A. Subjects and Experimental data

Seventeen University students (mean age: 20.58, stddev: 2.39 ; 10 men, 7 women) are voluntarily participated in this study. All subjects' native language were Turkish. They learned reading without any problem in the elementary school, however they were naive to the alphabet formed with distorted Turkish letters and the computer based training task. Before the experiment, all participants gave their informed consent after the experimental procedure was explained to them in accordance with guidelines set by the research ethics committee. Throughout the experiments, eMotiv EPOC+ headset is used. Internal sampling rate in the headset is 2048 per secs per channel. The data are filtered to remove mains artefacts then down-sampled to 128 per secs per channel. There are 14 EEG channels plus two references. Electrodes were placed according to the 10-20 system. Before the experiments, the calibration of the eMotiv Headset on the subject's scalp is done with the Xavier Control panel of eMotiv, each electrode is made sure to transfer EEG data with high quality (see Figure 1).

### B. Study Design

Participating subjects performed a letter print learning task under simultaneous EEG recordings. The goal of the task was learning the name of new letters which are distorted 180




Figure 1: The 10-20 numbering system of eMotiv EPOC electrodes.

degrees in the y-axis (See Figure 1). Before the experiments, the EEG signals in the resting, eyes open state are measured for each subjects for five minutes and the data is stored in .csv files. During these recordings, subjects were placed approximately 0.5 meters in front of a computer screen and instructed to relax with eyes open, looking at a fixation point in the middle of the screen. The programs are written with Python and uses the Community SDK provided by eMotiv to communicate with eMotiv EPOC+ headset. Artefact removal and conversion from Analog to Digital signal is done by the eMotiv standard procedures. libEDK.IEE\_GetAverageBandPowers routine is used. During the experiments EEG is recorded, and after the experiment, another five minutes of resting, "eyes open" state EEG signals are recorded.

### C. Letter print learning task

The task involved showing distorted letters on the screen together with a picture which starts with the letter shown and the phonics of letter (See Figure 2). Throughout the experiments, subjects are shown 29 distorted Turkish letters, repeated three times. In each repetition, there are slight differences in the order of pictures and sounds which are shown on the screen. In the first round, the picture and sound are shown at the same time, in the second round, the picture is shown one second before the sound, in the third round, the picture is shown one second after the sound. This is done for addressing differences in the cross modal attention shift characteristics of different subjects.




Figure 2: The computer based training program to teach distorted letters.



TABLE I: Learning performance measures

ID	Age	Sex	error pre-training	error post training	Oneminute-pre	Oneminute-post	Words read one minute
307	19	Male	3	1	14	40	28
304	19	Female	3	2	49	75	28
5000	20	Male	4	1	42	65	26
5005	18	Female	7	1	18	31	18
5002	20	Female	6	5	31	46	16
302	22	Female	4	2	24	36	14
5006	20	Male	0	2	0	16	14
5004	19	Male	4	2	25	36	13
2002	20	Male	4	1	31	40	12
5009	20	Male	3	0	31	40	12
5001	20	Female	7	5	42	51	11
308	18	Male	0	0	30	38	8
303	27	Female	3	2	49	56	8
5003	20	Male	3	3	45	52	7
2001	25	Male	1	0	37	40	1
306	20	Male	1	0	50	43	-6
301	23	Female	2	2	43	32	-9

TABLE II: Theta powers at Broca area

ID	Words read in one minute	Avg Theta (F7)	Avg Theta(FC5)	Average of Theta at F7 and FC5
307	28	1,346621	1,564475	1,455548
304	28	0,821042	0,352808	0,586925
5000	26	1,233911	0,860058	1,0469845
5005	18	0,75083	1,582611	1,1667205
5002	16	2,077955	0,746231	1,412093
302	14	0,440884	2,090648	1,265766
5006	14	2,873486	1,878642	2,376064
5004	13	1,463205	0,562709	1,012957
2002	12	1,038742	1,374948	1,206845
5009	12	2,15392	0,934102	1,544011
5001	11	3,144928	2,957646	3,051287
308	8	1,571115	2,060929	1,816022
303	8	2,549321	2,086834	2,3180775
5003	7	3,630647	1,467857	2,549252
2001	1	0,915747	2,660747	1,788247
306	-6	3,537321	1,495707	2,516514
301	-9	2,6577	2,307136	2,482418

REFERENCES

[1] Torgesen, Joseph K., et al. "Contributions of phonological awareness and rapid automatic naming ability to the growth of word-reading skills in second- to fifth-grade children." *Scientific studies of reading* 1.2 (1997): 161-185.

[2] Geva, Esther, and Linda S. Siegel. "Orthographic and cognitive factors in the concurrent development of basic reading skills in two languages." *Reading and Writing* 12.1 (2000): 1-30.

[3] Dickinson, David K., et al. "The comprehensive language approach to early literacy: The interrelationships among vocabulary, phonological sensitivity, and print knowledge among preschool-aged children." *Journal of Educational Psychology* 95.3 (2003): 465.

[4] Siegel, Linda S. "IQ is irrelevant to the definition of learning disabilities." *Journal of learning disabilities* 22.8 (1989): 469-478.

[5] Spironelli, Chiara, Barbara Penolazzi, and Alessandro Angrilli. "Dysfunctional hemispheric asymmetry of theta and beta EEG activity during linguistic tasks in developmental dyslexia." *Biological psychology* 77.2 (2008): 123-131.

[6] Duffy, Frank H., et al. "Dyslexia: Regional differences in brain electrical activity by topographic mapping." *Annals of neurology* 7.5 (1980): 412-420.

[7] Hallahan, Daniel P. "Some thoughts on why the prevalence of learning disabilities has increased." *Journal of Learning Disabilities* 25.8 (1992): 523-528.

[8] Ekanayake, H. (2010). "P300 and Emotiv EPOC: Does Emotiv EPOC capture real EEG?" Retrieved from: <http://neurofeedback.visaduma.info/emotivresearch.htm>

[9] Ozdenizci, Ozan, et al. "Electroencephalographic identifiers of motor adaptation learning." *Journal of Neural Engineering* (2017).

[10] Gitelman, Darren R., et al. "Language network specializations: an analysis with parallel task designs and functional magnetic resonance imaging." *Neuroimage* 26.4 (2005): 975-985.

[11] D'Angiulli, Amedeo, and Linda S. Siegel. "Cognitive functioning as measured by the WISC-R: Do children with learning disabilities have distinctive patterns of performance?." *Journal of Learning Disabilities* 36.1 (2003): 48-58.

[12] Ross, Gail, et al. "Effects of mothers' autoimmune disease during pregnancy on learning disabilities and hand preference in their children." *Archives of pediatrics and adolescent medicine* 157.4 (2003): 397-402.

[13] Wood, Lawrence C., and David S. Cooper. "Autoimmune thyroid disease, left-handedness, and developmental dyslexia." *Psychoneuroendocrinology* 17.1 (1992): 95-99.

[14] KORKMAZLAR-ORAL, Umran. "OGRENME SORUNLARINI ERKEN TANIMAK." *CEREBRAL PALSY SYMPOSIUM V*.

[15] Korkmazlar, Ü. "6-11 yas ilkokul cocuklarinda ozel ogrenme bozuklugu ve tani yontemleri" *Doktora Tezi, İstanbul Üniversitesi Tıp Fakültesi, İstanbul* (1992).

[16] Karaman, D., T. Turkbay, and F. S. Gokce. "Ozgul ogrenme bozuklugu ve dikkat eksikligi hiperaktivite bozuklugu binisikliginin bilissel ozellikleri." *Cocuk ve Genclik Ruh Sagligi Dergisi* 13.2 (2006): 60-8.

[17] KARAMAN, Dursun, Koray KARA, and İbrahim DURUKAN. "OZGUL OGRENME BOZUKLUGU." *Anatolian Journal of Clinical Investigation* 6.4 (2012).

[18] Bamidis, Panagiotis, John Gruzelier, and Ana Vivas. "SAN 2011 MEETING PROGRAM."

[19] Liu, Yue, et al. "Implementation of SSVEP based BCI with Emotiv EPOC." *Virtual Environments Human-Computer Interfaces and Measurement Systems (VECIMS), 2012 IEEE International Conference on. IEEE, 2012.*

[20] Mayaud, Louis, et al. "A comparison of recording modalities of P300 event-related potentials (ERP) for brain-computer interface (BCI) paradigm." *Neurophysiologie Clinique/Clinical Neurophysiology* 43.4 (2013): 217-227. bibitemBou Boutani, Hidenori, and Mieko Ohsuga. "Applicability of the "Emotiv EEG Neuroheadset" as a user-friendly input interface." *Engineering in Medicine and Biology Society (EMBC), 2013 35th Annual International Conference of the IEEE. IEEE, 2013.*

[21] Papagiannopoulou EA, Lagopoulos J. Resting State EEG Hemispheric Power Asymmetry in Children with Dyslexia. *Frontiers in Pediatrics*. 2016;4:11. doi:10.3389/fped.2016.00011.

[22] Wijers, Albertus A., Pieter H. Been, and Koos S. Romkes. "Dyslexics show a deviant lateralization of attentional control: a brain potential study." *Neuroscience letters* 374.2 (2005): 87-91.

[23] Hugdahl, Kenneth, and Richard J. Davidson, eds. *The asymmetrical brain*. MIT press, 2004.

[24] Kang, D.-H., Davidson, R. J., Coe, C. L., Wheeler, R. E., Tomarken, A. J., Ershler, W. B. (1991). Frontal brain asymmetry and immune function. *Behavioral Neuroscience*, 105(6), 860-869.

[25] Sangal, R. Bart, and JoAnne M. Sangal. "Use of EEG beta-1 power and theta/beta ratio over Broca's area to confirm diagnosis of attention deficit/hyperactivity disorder in children." *Clinical EEG and neuroscience* 46.3 (2015): 177-182.

[26] Lubar, Judith O., and Joel F. Lubar. "Electroencephalographic biofeedback of SMR and beta for treatment of attention deficit disorders in a clinical setting." *Applied Psychophysiology and Biofeedback* 9.1 (1984): 1-23.

[27] Saloni Krishnan, Kate E.Watkins, Dorothy V.M.Bishop . "Neurobiological Basis of Language Learning Difficulties" *CellPress Volume 20, Issue 9* (2016).